

Training Module #2

1. All are appropriate circumstances for perinatal depression screening except
 - a. Well-visit, 2-month-old accompanied by mother
 - b. Well-visit, 1-month-old accompanied by foster mother to whom she was discharged from NICU
 - c. Well-visit, 1 month old accompanied by grandmother because mother has a viral illness and is at home
 - d. Well-visit, 4-month-old accompanied by father
 - e. Well-visit, 6-month-old accompanied by mother identified with depression at 2- month visit

2. Which is a component of primary care intervention after screening? (check all that apply)
 - a. Demystification
 - b. Encouragement of social connections
 - c. Encourage turning over care of the child to another family member
 - d. Use of Common elements
 - e. Support for breastfeeding

3. Which are social-emotional screening tools for infants and young children? (check all that apply)
 - a. BITSEA (Brief Infant Toddler Social Emotional Assessment)
 - b. PEDS (Parents' Evaluation of Developmental Status)
 - c. BPSC (Baby Pediatric Symptom Checklist)
 - d. ASQ:SE -2 (Ages & Stages Questionnaire: Social-emotional -2)
 - e. PPSC (Preschool Pediatric Symptom Checklist)

4. A 2-way release is required for communication with all except:
 - a. Part C Early Intervention
 - b. Mental health clinician for the infant/dyad
 - c. Primary Care Clinician of the mother
 - d. Mental health clinician for the mother

5. What advocacy issues are relevant to your region/state? (check all that apply)
- a. Expansion of Medicaid benefits for pregnant women beyond the postpartum period
 - b. Expansion of the mental health workforce for infant and early child mental health (IECMH) professionals
 - c. Adequate payment for mental health providers serving infants and young children
 - d. Expansion of Early Intervention eligibility for risk to social-emotional development
 - e. Expansion of mental health services for women with perinatal depression



Answer Key:

1. c
2. a, b, d, e
3. a, c, d, e
4. b
5. a, b, c, d, e